Ophthalmology and Visual Sciences



Bilateral Dacryocystoceles with Acute Dacryocystitis

Category(ies): Oculoplastics, Pediatrics

Contributor: <u>Lucas Lenci, MD</u> Photographer: Lucas Lenci, MD



6-day-old with worsening redness around the right eye. These images provide an example of acute dacryocystitis in the setting of bilateral dacryocystoceles.



This patient's left side is a classic example of a dacryocystocele which typically presents as a gray to bluish mass or region of swelling inferior to the medial canthal tendon. An area of swelling above this level is worrisome for conditions such as encephalocele or dermoid tumors. Patients need to be monitored closely as secondary infections can develop.





The patient has developed acute dacryocystitis on the right. Classically, dacryocystitis presents with redness, swelling, pain, and purulent discharge on the affected side. This patient was admitted to the hospital for IV antibiotic therapy followed by probing of the nasolacrimal system.





Image Permissions:



Ophthalmic Atlas Images by <u>EyeRounds.org</u>, <u>The University of Iowa</u> are licensed under a <u>Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License</u>.

Address

University of Iowa Roy J. and Lucille A. Carver College of Medicine Department of Ophthalmology and Visual Sciences 200 Hawkins Drive Iowa City, IA 52242

Support Us

Legal

Copyright © 2019 The University of Iowa. All Rights Reserved Report an issue with this page Web Privacy Policy | Nondiscrimination Statement

Related Links

Cataract Surgery for Greenhorns EyeTransillumination Gonioscopy.org Iowa Glaucoma Curriculum Iowa Wet Lab Patient Information Stone Rounds The Best Hits Bookshelf

EyeRounds Social Media

Follow







Receive notification of new cases, sign up here Contact Us Submit a Suggestion