

Myasthenia Gravis - edrophonium test

Category(ies): Neuro-ophthalmology

Contributor: [Jordan Graff, MD](#)



The materials needed for an edrophonium (Tensilon, Enlon) test for myasthenia gravis are here pictured. Atropine may be used to prevent cardiac side effects of edrophonium. Serum can be sent for Acetylcholine Receptor antibodies.



EyeRounds.org

Pupils were equal and reactive with a >3 log unit RAPD OD. Dilated funduscopic examination showed severe, diffuse optic atrophy OD and bow-tie (band) optic atrophy OS.

 [Enlarge](#)  [Download](#)

Image Permissions:



Ophthalmic Atlas Images by [EyeRounds.org](#), [The University of Iowa](#) are licensed under a [Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License](#).

Address

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences

Legal

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)

Related Links

Cataract Surgery for Greenhorns
EyeTransillumination
Gonioscopy.org
Iowa Glaucoma Curriculum

EyeRounds Social Media

Follow



200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

[Receive notification of new cases,
sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)