

Prostaglandin analogue-induced orbital fat atrophy

Category(ies): Glaucoma, Iris, Oculoplastics

Contributor: Jeff Welder, MD



52-year-old female with angle recession glaucoma of the left eye treated with a prostaglandin analogue for many years with resultant orbital fat atrophy on that side. Note the superior sulcus.

[Enlarge](#) [Download](#)



The patient returned one year after discontinuing the prostaglandin analogue and her orbital fat atrophy had markedly improved.

[Enlarge](#) [Download](#)



Ophthalmic Atlas Images by [EyeRounds.org](https://www.eyerounds.org), [The University of Iowa](https://www.theuniversityofiowa.edu) are licensed under a [Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License](https://creativecommons.org/licenses/by-nc-nd/3.0/).

Address

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences
200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

Legal

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)
[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

Related Links

[Cataract Surgery for Greenhorns](#)
[EyeTransillumination](#)
[Gonioscopy.org](#)
[Iowa Glaucoma Curriculum](#)
[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

EyeRounds Social Media

Follow



Receive notification of new cases,
[sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)